


Accepted Manuscript

Transcriptome analysis of the synganglion from the honey bee mite, *Varroa destructor* and RNAi knockdown of neural peptide targets

Ewan M. Campbell, Giles E. Budge, Max Watkins, Dr Alan S. Bowman



PII: S0965-1748(15)30081-3

DOI: [10.1016/j.ibmb.2015.12.007](https://doi.org/10.1016/j.ibmb.2015.12.007)

Reference: IB 2801

To appear in: *Insect Biochemistry and Molecular Biology*


Received Date: 4 August 2015

Revised Date: 4 December 2015

Accepted Date: 20 December 2015

Please cite this article as: Campbell, E.M, Budge, G.E, Watkins, M., Bowman, A.S, Transcriptome analysis of the synganglion from the honey bee mite, *Varroa destructor* and RNAi knockdown of neural peptide targets, *Insect Biochemistry and Molecular Biology* (2016), doi: 10.1016/j.ibmb.2015.12.007.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Transcriptome analysis of the synganglion from the honey bee mite, *Varroa destructor* and RNAi knockdown of neural peptide targets.

Authors: Ewan M Campbell^a, Giles E Budge^b, Max Watkins^c, Alan S Bowman^a

^a School of Biological Sciences (Zoology), University of Aberdeen, Tillydrone Avenue,
Aberdeen AB24 2TZ, UK

^b National Bee Unit, Fera, Sand Hutton, York, YO41 1LZ, UK

^c Vita (Europe) Limited, Vita House, London Street, Basingstoke, Hampshire, RG21 7PG
UK

Corresponding Author:

Dr Alan S. Bowman
School of Biological Sciences (Zoology)
University of Aberdeen
Aberdeen
AB24 2TZ

Tel: +44 1224 272877
Fax: +44 1224 272396
Email: a.bowman@abdn.ac.uk

Abstract:

Varroa mites (*Varroa destructor*) and the viruses that they transmit are one of the major contributing factors to the global honey bee crisis. Gene products within the nervous system are the targets of all the insecticides currently used to control *Varroa* but there is a paucity of transcriptomic data available for *Varroa* neural tissues. A cDNA library from the synganglia (“brains”) of adult female *Varroa* was constructed and 600 ESTs sequenced and analysed revealing several current and potential druggable targets. Contigs coding for the deformed wing virus (DWV) variants *Varroa destructor* virus-1 (VDV-1) and the recombinant (VDV-1_{DVD}) were present in the synganglion library. Negative-sense RNA-specific PCR indicated that VDV-1 replicates in the *Varroa* synganglion and all other tissues tested, but we could not detect DWV replicating in any *Varroa* tissue. Two neuropeptides were identified in the synganglion EST library: a B-type allatostatin and a member of the crustacean hyperglycaemic hormone (CHH) superfamily. Knockdown of the allatostatin or the CHH-like gene by double-stranded RNA-interference (dsRNAi) resulted in 85% and 55% mortality, respectively, of *Varroa*. Here, we present the first transcriptomic survey in *Varroa* and demonstrate that neural genes can be targeted by dsRNAi either for genetic validation of putative targets during drug discovery programmes or as a potential control measure in itself.

Keywords:

Varroa destructor, *Apis mellifera*, synganglion, deformed wing virus, allatostatin, crustacean hyperglycaemic hormone, gene knockdown

1. Introduction

The economic contribution of insect pollination to crop production is significant (Gallai et al., 2009) and managed honey bees provide about half of this ecosystem service to all insect-pollinated crops on Earth (Kleijn et al., 2015). The critical importance of honey bees (*Apis mellifera*) to global food production, and the central role of *Varroa destructor* to *A. mellifera* colony losses, elevate *Varroa* to arguably the most serious parasite of any livestock species worldwide. *V. destructor* is an ectoparasitic mite of honey bees that is now endemic in all countries where husbandry of honey bees occurs, except Australia (vanEngelsdorp et al., 2009, Martin et al., 2013, vanEngelsdorp and Meixner, 2010). *V. destructor* are vectors of at least 18 different honey bee viruses, including highly pathogenic strains of deformed wing virus (DWV) (Di Prisco et al., 2011, Wang et al., 2013, Genersch and Aubert, 2010). The recent spread of *Varroa* coupled with its ability to act as a vector for DWV has resulted in a synergistic effect that has been heavily implicated in a global decline in honey bee health (vanEngelsdorp et al., 2009, Martin et al., 2012). *Varroa* and associated pathogens have led to higher than average winter losses in many regions (Highfield et al., 2009), resulting in increased effort and money spent on combating infestations and replacing stock. Typically, hives infested with *Varroa* are unlikely to survive without intervention for more than 1-3 years (Martin, 1998). Integrated pest management strategies including drone brood culling, mite population monitoring and the use of various organic acids during broodless periods can significantly reduce *Varroa* burden (Rosenkranz et al., 2010). Breeding *Varroa* resistant or tolerant *A. mellifera* offers great potential, but, is not yet a reality (Correa-Marques et al., 2002). The control of *Varroa* still remains largely based on chemical acaricides. There is now widespread and increasing resistance to tau-fluvalinate, flumethrin, coumaphos and amitraz (Rodriguez-Dehaibes et al., 2011), leaving many areas with no effective control measures against *Varroa* during much of the beekeeping season. The current lack of alternative control methods may undermine the future of sustainable apiculture globally (Dietemann et al., 2012) and, hence, general food security.

In common with other pesticides, the majority of acaricides target vital pathways and channels within the nervous system (Lees et al., 2010) and it is highly probable that the development of new acaricides will also target this tissue. Sequencing of the *V. destructor* genome and tissue-specific transcriptome libraries, along with reverse genetic approaches such as gene-knockdown by RNA interference (RNAi) can help to identify physiologically important pathways and inform selection of viable targets for therapeutic control. Given the

importance of *Varroa* to global food production, there are rather limited genetic resources other than an initial genomic survey (Cornman et al., 2010). Presently, there is an absence of any neural-tissue specific *Varroa* sequences in public databases other than the voltage-gated sodium channel (Wang et al., 2003), the target of pyrethroids.

RNAi is used widely as a tool for investigating gene function in many taxa, including arachnids (Karim et al., 2008, Browning and Karim, 2013), and has been employed to investigate the physiology as well as inform acaricide development in numerous tick species. RNAi is effective in the two-spotted spider mite *T. urtica* (Khila and Grbic, 2007) and recently was demonstrated to be effective in *V. destructor* directly, through injection and immersion of dsRNA (Campbell et al., 2010) or via feeding dsRNA to honey bees on which *Varroa* are present (Garbian et al., 2012). To date, RNAi in *Varroa* has been limited to targeting multi-tissue housekeeping transcripts, such as V-ATPase subunits and others involved with cellular homeostasis or detoxifying enzymes ubiquitously expressed in all tissues. RNAi has huge potential for applied entomology (Scott et al., 2013, Price and Gatehouse, 2008). Neural targets for the development of pesticides can be tested by high throughput RNAi gene-knockdown (Lees and Bowman, 2007) and it has become increasingly apparent that RNAi has the potential itself to manage pests and parasites by targetting transcripts that are vital to pest physiology by suppressing vital genes to reduce reproductive capacity and increase mortality (Scott et al., 2013). It is reasonable to postulate that the synganglion is the site of numerous genes highly critical to the *Varroa*'s survival, but the sequence of such genes is not currently available.

In the current study, we constructed and sequenced an EST library from *Varroa* synganglia to identify neural genes that may be potential targets for *Varroa* control. We then investigated the potential for dsRNA to silence two neurohormone genes identified within the *Varroa* synganglion library. Both targets were inhibited by dsRNA and displayed individual phenotypic responses to RNAi including increased mortality. This work demonstrates dsRNA can gain access to the synganglion of *Varroa* enabling further investigation into mite neurophysiology and inform future methods of control. The lethality of neural peptide gene knockdown also heralds the possibility of *Varroa* control by RNAi of neural tissue –specific targets.

2. Materials and methods

2.1. EST library construction

Varroa destructor (adult female) mites were collected from capped brood cells frames from *Apis mellifera* hives from York, England that had purposefully been left untreated for *Varroa* control. Prior to harvesting mites, the bee frames were maintained at 27°C and 80% relative humidity environment with a 15.5 h : 8.5 h light:dark regime. Approximately 50 synganglia were dissected from mites in phosphate buffered saline (PBS) before being washed in sterile ice-cold PBS and pooled together in a 1.5 ml eppendorf microfuge tube containing 200 µl RNA-later (Sigma, Poole, UK). De-synganglionized mite bodies as well as Malpighian tubules, gut and whole mites were also retained. Prior to RNA extraction, an additional 450 µl dissection buffer was added to sample tubes and centrifuged at 20,000 RCF for 15 min at 4°C. Supernatant was removed and the synganglion washed with fresh PBS before a final centrifuge again at 20,000 RCF for 15 min at 4°C. Supernatant was again removed and 600 µl ZR extraction buffer added to each tissue sample. Total RNA was extracted using a mini-RNA isolation II Kit (Zymo Research, Orange, California, USA), as per manufacturer's instructions and eluted in 50 µl water. RNA was co-precipitated with 1.5 µl glycogen blue (Life Technologies, Paisley UK) and 5 µl 3M sodium acetate in 95% ethanol and resuspended in 5 µl of DEPC-treated water.

First strand cDNA synthesis was completed using 3.5 µl (0.5 µg) of total RNA. The construction of cDNA libraries was done using the SMART cDNA library construction kit (Clontech, St-Germain-en-Laye, France) according to the protocol provided by manufacturer, with some modifications. To determine optimal number of cycles, two identical amplification reactions were prepared. After the 10th amplification cycle the first reaction was stored on the ice, while the second one was used for the PCR cycle number optimization by removing 3 µl samples from the reaction tube every two cycles until cycle number 20. Samples were checked by visualization on a 1.1% agarose gel. The optimal number of cycles with visible and equally represented products, in this case 20 cycles, was used for primary amplification. cDNA was proteinase K treated extracted using phenol:chloroform and resuspension in water. After *Sfi*I digestion and size fractionation with a Chroma Spin-400 column, fractions were checked using agarose gel and pooled into large or medium libraries. Pooled cDNA was ethanol precipitated and eluted in 4 µl of water prior to 3 µl from each fraction being ligated into the λTripleEx2 vector and packed into phage using the Gigapack III Gold Packaging extract (Stratagene). Each un-amplified library was mixed with *E.coli* XL1 blue cells and top

agar supplemented with X-gal and IPTG before being plated onto LB MgSO₄ agar plates in serial dilutions of neat, 1:10, 1:100 and 1:1000.

In total, 600 randomly selected recombinant plaques (white) were picked as agar plugs into 96-well plates, each well containing 100 µl of SM buffer (0.58% NaCl, 0.2% MgSO₄·H₂O, 0.05M Tris-HCl, pH 7.5, 0.02% gelatin). Four plates (384 clones) were picked from the large fraction library, two plates (192 clones) from the medium fraction library and an additional 24 clones from the large fraction library for initial quality control. PCR with vector-specific primers was carried out using SM buffer / picked plaques as templates. PCR was carried out in 96-well plates containing 25 µl 2x Biomix (Bioline), 5 µl template, 1 µl (10 ng/µl) each of PT2F1 (5'- AAGTACTCTAGCAATTGTGAGC-3') and PT2R1 (5'- CTCTTCGCTATTACGCCAGCTG- 3') and 18 µl water to give a 50 µl final reaction volume. Cycling conditions were 94°C for 15 min followed by 33 cycles of 94°C for 1 min, 49°C for 1 min and 72°C for 1 min 20s. PCR products were sent to GATC (Konstanz, Germany) for PCR reaction clean up and sequenced using primer PT2F3 (5' – CTCGGAAGCGCGCCATTGT- 3'). PT2F3 is upstream from inserted cDNA and downstream from PT2F1 primer used in initial PCR reaction.

2.2. Bioinformatics

Expressed sequence tags (ESTs) were trimmed of primer and vector sequences, clustered and checked for sequence quality using Lasergene Seqman (LaserGene v8.03, DNASTar, Madison, USA). BLASTn, BLASTx and tBLASTx programmes were used within the programme BLAST2GO to compare the EST nucleotide sequences with the nonredundant (NR) databases of the NCBI and to the Gene Ontology (GO) database (www.blast2go.org). BLASTn against the *Varroa* WGS genome was carried out using the NCBI BLAST servers. Following analysis of results, transcripts were initially classified as novel sequences, putative identity or unknown function. Transcripts with a putative identity were further divided into functional categories by analysing GO identity and homology to known genes.

Neurohormone peptides were identified by signal peptide sequence, pro-hormone processing sites and length (<300 amino acids). Signal peptide analysis was carried out using SignalP (<http://www.cbs.dtu.dk/services/SignalP/>) and Signal-3L (<http://www.csbio.sjtu.edu.cn/bioinf/Signal-3L/>) using both the neural networks and Hidden Markov Model algorithms. Pro-hormone cleavage sites were predicted based on work by

Veenstra (Veenstra, 2000), and Neuropred (<http://neuroproteomics.scs.illinois.edu/neuropred.html>). Predicted transmembrane topology of the neurohormones was predicted with TMHMM (Krogh et al., 2001) at <http://www.cbs.dtu.dk/services/TMHMM/>.

The neurohormones crustacean hyperglycaemic hormone (CHH) and allatostatin (AST) sequences were primarily aligned in GUIDANCE using the progressive algorithms PRANK and MAAFT (<http://guidance.tau.ac.il/>) with confidence scores assigned for each residue, column, and sequence. Unreliable columns and sequences were removed from the initial alignment in preparation for downstream analyses. MEGA 6.3 (Tamura et al., 2013) was used to align sequences by MUSCLE. Phylogenetic trees were constructed by Maximum likelihood using the LG (+G+I) model of analysis and bootstrapped 1000 times.

2.3. Virus identity, distribution and replication

Three sequences with homology to *Varroa destructor* virus-1 (VDV-1) were found in the EST library as determined by BLASTn. Alignment comparisons were carried out with MEGA(6.3) using genome data from known VDV-1 and DWV strains (Moore et al., 2011, de Miranda et al., 2013). Pairwise identity comparison was determined by CLUSTAL-omega.

To determine the variety of DWV strains present in the synganglia of *Varroa*, as well as tissue specificity, a nested-PCR approach was taken. Generic primers spanning the major recombination junction between DWV and VDV-1 were used in the first round PCR (Moore et al., 2011) using synganglia EST library material as well as cDNA constructed from gut, Malpighian tubules and de-synganglionized mite carcass. Nested-PCR was then carried out with variant specific primers and 1 µl of first-round product. Products were visualized by gel electrophoresis.

Virus replication was assessed by the specific detection of negative strand viral RNA using strand-specific Reverse Transcriptase-PCR (Yue and Genersch, 2005). 0.5 µg of mite RNA extracted from synganglia, gut, Malpighian tubules and de-synganglionized mite carcass (see above) was reverse transcribed with thermostable Superscript III (Life Technologies, Paisley, UK) and tagged with first-strand primers specific for either DWV or the VDV variant of the virus. Reverse transcription was carried out at 55°C for 1 hour and treated with RNase-H to remove the possibility of false positive detection during subsequent PCR. A template-free cDNA reaction was carried out as a control. Tissue specific virus-primed cDNA was

subjected to qPCR by SybrGreen detection on an Opticon II platform PCR cycling conditions were as follows: 1 cycle of 10 min at 95°C, followed by 35 cycles of 20s at 94°C, 20s at 55°C and 20s at 72°C. Control reactions with primer and template free reaction mixture were included. To confirm product specificity, melting point analysis was performed at the end of the qPCR run and the products analysed by gel electrophoresis.

2.4. Spatial distribution of neural peptides

To determine if CHH and AST transcripts are restricted to the synganglionic tissue of *Varroa*, RNA extracted from whole mites, de-synganglionized mites, gut tissue and Malpighian tubules was reverse transcribed. After isolation, 0.5 µg total RNA was DNase treated with 1 µl (2U) RQ1-DNase (Promega, Southampton, UK) and 1 µl RQ1 buffer and incubated at 37°C for 30 min. DNase-treated total RNA was incubated at 70°C with 0.5 µg of oligo d(T)15 (Promega) in a total volume of 10 µl for 5 min. Material was snap-chilled on ice for 5 min prior to the addition of 5 µl 5×RT buffer, 1 µl dNTPs (25 mM each), 0.5 µl Bioscript-reverse transcriptase and DEPC water to 25 µl. The reaction was incubated at 42°C for 1 hour. PCR was carried out with CHH, AST and *Varroa* actin specific primers using the following cycling conditions: 1 cycle of 5 min at 94°C, followed by 35 cycles of 1 min at 94°C, 1 min at 53°C and 45 s at 72°C. Products were resolved on an agarose gel and visualised by UV light.

2.5. Preparation of dsRNA

VdCHH and *VdAST* dsRNA was prepared with BLOCK-iT RNAi TOPO transcription kits (Invitrogen, Paisley, UK), according to the manufacturer's instructions. LacZ-dsRNA was prepared and used as a negative control. Briefly, PCR was carried out using adult female *V. destructor* cDNA in conjunction with *VdCHH* specific primers (*VdCHH* F1/R1), *VdAST* specific primers (*VdAST* F1/R1) or with control LacZ-plasmid and LacZ specific primers (LacZ-F2, ACCAGAAGCGGTGCCGAAA and LacZ-R2, CCACAGCGGTGGTTCGGAT). PCR cycling conditions were as described above followed by a final extension time of 15 min at 72°C. Products were resolved on an agarose gel, excised and purified using a Qiagen gel extraction kit (Qiagen, Crawley, UK). TOPO-T7 linker was ligated to *VdAST*, *VdCHH* and LacZ reactions before a secondary PCR was carried out to produce sense and antisense templates. T7-RNA polymerase was used in transcription reactions with target templates to generate sense and antisense RNA. Single RNA strands were annealed and the resultant dsRNA purified and quantified by a ND-1000

Nanodrop Spectrophotometer (Labtech Inc., East Sussex, UK). dsRNA was ethanol precipitated and resuspended in DEPC-treated water to a working concentration of 2.5 µg/µl and stored at -80°C.

2.6. Administration of dsRNA by injection and immersion

Adult female *V. destructor* were removed from capped brood cells along with associated bee larvae. Microinjections were carried out according to methods previously described (Campbell et al., 2010). Briefly, mites were placed on double-sided tape ventral side up, and injected with 20 nl (2.5 µg/µl) of either *VdCHH*-dsRNA, *VdAST*-dsRNA or *LacZ*-dsRNA in the soft tissue either proximal to the anal region or postcoxal plate using pulled glass capillary needles in conjunction with a Harvard micro-injector system. Mites were placed in petri dishes containing 1 bee larvae per 4 mites. Dead or unhealthy looking mites, presumably due to the injection trauma, were discarded after 1 hour. To assess penetration of neural tissue by the immersion method (Campbell et al., 2010, Campbell et al., 2010), adult mites were removed from capped brood cells and placed in 500 µl microfuge tubes containing 20 µl *VdCHH*-dsRNA, *VdAST*-dsRNA or *LacZ*-dsRNA (2.5 µg/µl) in 0.9% NaCl. Mites were soaked at 4°C overnight, then removed, dried and placed in Petri dishes. Mites were fed on similar aged developing bee larvae (replaced every 24h) and maintained at 30°C and 85% RH.

2.7. Validation of RNAi

RNA was extracted from individual *Varroa* 48 h post-treatment. In addition, persistence of the RNAi effect was measured in injected mites by harvesting at 18, 24, 48 and 72 h. Total RNA was extracted from mites using ZR Insect RNA MicroPrep (Zymo Research, CA, USA), DNase-treated and reverse transcribed. PCR was carried out using either *VdCHH*-dsRNA, *VdAST*-dsRNA or *LacZ*-dsRNA treated sample cDNA in conjunction with primers specific for actin or *VdCHH* / *VdAST* using primers and cycling conditions, as described above. Products were visualized on an agarose gel normalized to actin loading. To assess persistence and approximate % knockdown, ImageJ software was used to carry out semi-quantitative densitometric analysis on gel images.

2.8. Phenotypic responses to RNAi

To assess mortality induced by neurohormone gene knockdown, adult female *Varroa* were removed from capped brood cells, treated with dsRNA by the immersion method described

above and maintained on developing bee larvae at 30°C and 85% RH. *Varroa* were monitored every 8 hours for up to 5 days and dead mites removed and counted. “Self righting” behaviour was assessed during inspections to determine morbidity and other non-typical behaviour was noted.

3. Results

3.1. Synganglion EST database summary

A *V. destructor* EST library derived from the synganglia of 50 mites’ RNA was constructed. The large fragment library consisted of 6.23×10^6 colony forming units (cfu)/ml and the medium fragment library consisted of 1.07×10^7 cfu/ml with recombination of 94.3 and 96.3%, respectively. A total of 600 plaques were picked from plates from the large and medium fractionated libraries and sequenced. 555 (92.5%) ESTs were of sufficient length (>75 bp) and quality for further analysis and assembled into 398 clusters, with a mean length of 484 bp, ranging from 92 – 899 bp. Of these clusters, 11 had more than one associated EST, with a mean of 15.2 ESTs / contig. The top three best coverage contigs were *V. destructor* mitochondrial genes; 16S ribosomal protein, 12S ribosomal protein and cytochrome oxidase containing 80, 28 and 10 ESTs, respectively. The remaining ESTs consisted of 387 singletons (Supplementary Table 1).

All clusters were initially compared by BLASTn and BLASTx to the *Varroa destructor* WGS (PRJNA33465) and mitochondrial (AJ493124.2) genomic databases. 381 clusters (95.7%) were homologous to database sequences. The remaining 17 clusters were BLASTed against other nr and protein databases using BLASTn and tBLASTx, respectively. A single cluster containing 3 ESTs from this subset was homologous to a region of *Varroa destructor* virus-1 (VDV-1), a pathogenic variant of DWV. Viral contigs were located near the 3’ end of the VDV-1 genome in a region coding for RNA-dependant RNA polymerase (RdRp). No classic DWV sequences as normally present in bees were isolated from the synganglia EST.

Clusters were BLASTed against NCBI nucleotide and protein collection (nr/nt) and assigned into categories (Fig. 1A). Of the 398 clusters analysed, 3.7% were novel and 33.4% had high homology to annotated sequences (Supplementary Table 1). The remaining clusters were homologous to existing *Varroa* genomic databases with no annotation or known identity. The 3.7% of sequences with no homology to the *Varroa* genome were BLASTed against other databases but no identity was determined. It may be that future versions of the *Varroa*

genome or on-going transcriptome studies will help with the identification of these contigs. Clusters with putative shared identity with existing sequences in the nr/nt database were compared with the GO database using BLAST2GO and assigned functional classification (Fig. 1B). The largest functional sets were for clusters associated with metabolism (24.7%) as well as cell growth, division or RNA transcription (20.3%). The potentially druggable target categories of transmembrane receptors and neurohormones accounted for 6.5% of synganglion library clusters. Sequence data and initial homology searches for all 398 clusters are given in Supplementary Table 1.

3.2. Virus identity, distribution and replication

Three contigs (L4C04, MIC05 and L1A04) coded for DWV variants. Contigs MIC05 and L1A04 were identical to each other and were treated in the analysis as such. L1A01 and L4A04, sequenced from the EST synganglia database both had >99% identity with VDV-1 (Genbank AY251269.2) but lower identity to DWV (NC_004830.2), 91% and 84%, respectively (Fig. 2a). To determine the DWV strain diversity in the synganglia of *V. destructor* we carried out strain-specific PCR across the identified recombination point. Nested PCR resulted in bands of the expected size for both VDV-1 as well as the recombinant VDV-1_{DVD} but DWV and recombinant VDV-1_{VVD} were absent (Fig. 2b).

To determine if VDV-1 is replicating in *Varroa* synganglia, we assayed for negative-sense RNA strands as during replication the positive-sense RNA of VDV-1 is transcribed to a full-length copy of the negative-sense RNA strand genome. Reverse transcription was carried out on RNA extracted from *Varroa destructor* tissues using VDV-1 or DWV specific primers and thermostable transcriptase. qPCR of VDV-1- or DWV-primed cDNA detected the presence of negative-strand RNA from VDV-1 in all tissues assayed as determined by gel electrophoresis (Fig. 2c) and product melting point analysis (data not shown). No negative-strand DWV was detected in any tissue assayed

3.3. Neuropeptide sequence analysis

3.3.1. CHH/GIH/MIH/CHH contig

A 705 bp contig found in the EST library (L1C01) was identified by BLAST to have high homology to crustacean hyperglycaemic hormone (CHH) from *Homarus americanus* (2105187B) and *Callinectes sapidus* (AAS45136) and to ion transport peptide (ITP) from the

tick *Dermacentor variabilis* (ACC99599.1). This sequence was identical to a region of the *Varroa* genome WGS (VDK00079049) and had high homology (88%) to a sequence found in the Western predatory mite, *Metaseiulus occidentalis*, genome (XM_003741077). No hits were returned when the coding region was BLASTed against the spider mite *Tetranychus urticae* WGS database.

In silico analysis of L1C01 contig demonstrated a coding region of 134 amino acids with a crustacean hyperglycemic hormone/molt-inhibiting hormone/gonad-inhibiting hormone (CHH/MIH/GIH) motif (Pfam PF011147) at positions 54 to 129. SignalP revealed a 28 amino acid signal peptide cleaved at position T28/Q29. The *M. occidentalis* sequence also contained the CHH/MIH/GIH motif and a signal peptide cleavage site at position A24/F25. Both sequences contain 6 canonical cysteine residues conserved in the CHH superfamily (Fig. 3c).

DELTA-BLASTp interrogation of the nr protein database with the L1C01 sequence returned multiple hits to members of the CHH-superfamily with six conserved cysteine residues identical across all results. Interestingly, most homology values are relatively low with maximum identities between 26 and 35% for sequences with significant expect values of 9e-05 and below. The top hit was an uncharacterized protein from the western predatory mite, *Metaseiulus occidentalis*. The top annotated hits are CHH sequences from various marine crustaceans whereas the highest ITP match was from the tick, *Dermacentor variabilis*. Top BLASTp hits and selected others mined from the *T. urticae* genome and tick databases were subjected to GUIDANCE analysis, aligned using MEGA6.3 and maximum-likelihood phylogenetic trees were constructed. Both *Varroa* and *M. occidentalis* sequences shared a branch along with other Acari sequences separated from both CHH, MIH/GIH and ITPs (Fig. 3). Although phylogenetic analysis determined the L1C01 contig resides within a distinct acarine clade, no inference of function was revealed and bootstrap values were relatively low between all clades.

3.3.2. Allatostatin contig

A 708 bp sequence found in the synganglia EST library (M2D11) was identified by BLAST to have homology (42%) to B-type allatostatin (AST) from *Aedes aegypti* (XP_001655873.1). This sequence was identical to a region of the *Varroa* genome WGS (VDK00000794) and had high homology (62%) to a sequence found in the Western

predatory mite, *Metaseiulus occidentalis*, genome (XP_003741714). *Varroa* AST belongs to the B-type AST as it contains the conserved peptide sequence motif W(X)6W and a weak signal peptide cleavage point between position A31 and V32 (Fig. 4a). The *M. occidentalis* AST also contains W(X)6W motifs and signal peptide cleavage point between positions A21 and K22. Similar to other arthropod B-type ASTs, *Varroa* AST has multiple lysine and arginine cleavage sites at positions 69/70, 87/88, 101/102, 112/113 and 127/128 that would result in three mature peptides 16, 12 and 13aa in length each containing the W(X)6W motif (Fig. 4b). Though three acarine B-type ASTs (*Varroa*, *M. occidentalis* and *I. scapularis*) contain three mature peptides, *T. urtica* contains six.

3.4. Neural peptide transcript distribution

RT-PCR demonstrated that AST is confined to the synganglion with no detection in Malpighian tubules, gut or de-synganglionized mite carcass. In contrast, though CHH was abundantly detected in the synganglion, it was lowly expressed in gut and the de-synganglionized carcass and very weakly present in Malpighian tubules (Fig. 5).

3.5. RNAi knockdown of neuropeptides

Successful knockdown for both AST and CHH was initially performed by intrahaemocoelic injection of the corresponding dsRNAi (data not shown). All subsequent knockdown experiments and presented data administered the dsRNA by immersion which allows more *Varroa* to be treated and decreases mortality from the injection trauma. *Varroa* treated with dsRNA-AST showed a decrease in AST mRNA levels within 24 hr and decreasing further by 85% knockdown ($P < 0.001$) of target gene after 48h compared to the control *Varroa* group treated with dsRNA- LacZ (Fig. 6a). Mites treated with dsRNA-AST exhibited significantly higher mortality from 24h ($P < 0.01$) onwards compared to controls with only 30% of mites surviving at 72 h ($P < 0.001$) (Fig. 6b). In repeated independent experiments (not shown), the level of mortality in dsRNA-AST treated *Varroa* was consistently 50 – 60% higher than the dsRNA-LacZ treated *Varroa* over similar time spans. Though AST-knockdown increased *Varroa* mortality, no obvious phenotypic changes were observed.

The level of gene knockdown for mites treated with dsRNA-CHH was ~55% after 48h compared to the dsRNA-LacZ treated controls ($P < 0.03$) (Fig. 7a). Mortality levels in *Varroa* treated with dsRNA-CHH were greater compared to dsRNA-LacZ treated mites attaining statistical significance after 72 h ($P < 0.05$). Survival after 96 h was 68% compared to 90%

in control groups ($P < 0.05$) (Fig. 7b). In a cohort of *Varroa* treated with dsRNA-CHH a notable “shaking” or “shivering” phenotype was observed in which the mite’s whole body shook from side to side while still on their feet and occurred whether the mite was on the bee larva or the Petri dish. This “shaking” was observed in all repeated experiments using different preparations of dsRNA-CHH and mites from different hives and was not observed in any mites treated with dsRNA-LacZ or dsRNA-AST. Because the *Varroa* were in groups and not individually marked, we could not determine if the “shaking” phenotype preceded death or if those mites recovered, but it was noted that groups of mites in which we observed “shaking” mites tended to have higher numbers of dead mites at the subsequent observation time point.

4. Discussion

Hormones and neurotransmitters play key roles in development and physiology (Kwon et al., 2001, Friis et al., 2004, Cabrera et al., 2009) and this paper presents the first transcriptome from *Varroa* and the first synganglion transcriptome in any mite. Information about neural targets, such as membrane channels, hormones and receptors provide an opportunity in the development of chemicals to control *Varroa* (Wolstenholme et al., 2007) while the RNAi gene knockdown approach described in this paper could be employed for rapid genetic validation of a given target ahead of the long chemical validation process.

Next-generation sequencing technologies are developing rapidly and are revolutionising genomics and transcriptomics but Sanger DNA sequencing techniques, such as presented here, still have an important role to play in the identification of physiological pathways that may be key to finding targets for pest control (Schwarz et al., 2013). Indeed even from the modest EST data presented here, we identified two hormones and multiple transmembrane receptors that are potential control targets, including a nicotinic acetylcholine receptor subunit, the target of neonicotinoids. The majority of contigs we sequenced mapped directly onto the first draft of the *Varroa* genome and 42.5% had putative identity to existing genes. However, this leaves 57.5% that still have no known function. Indeed 15 transcripts, or 3.7%, were not evident in the *Varroa* genome. This paucity of information highlights the limited molecular resources available to *Varroa* biologists and although the number of transcriptomic projects from closely related mites such as *Tetranychus urtica* (Bryon et al., 2013) and *M. occidentalis* (Hoy et al., 2013) is increasing, further work on annotating and characterising the *Varroa* genome and available transcriptome data is needed.

Within the *Varroa* synganglion library we identified relatively high levels (3/555 clones) of a pathogenic virus transmitted from *Varroa* to bees. DWV is almost ubiquitous in honey bee colonies and its transmission and association with *Varroa* increases significantly its pathogenicity to honey bees (Genersch and Aubert, 2010, Ryabov et al., 2014). DWV is closely related to VDV-1, with 84% nucleotide (95% amino acid) identity (Ongus et al., 2004) and this clade of viruses contribute significantly to the collapse of honey bee colonies. Sequence analysis of EST transcripts from the *Varroa* synganglion EST revealed the presence of three contigs with nearly 100% identity to *Varroa destructor* virus-1, spanning the 3' end of the viral sequence but no DWV transcripts were found. RT-PCR using virus specific primers confirmed the presence of VDV-1 in the synganglion library and revealed it was present throughout the gut, Malpighian tubules and in whole mites demonstrating no dramatic tropism for a particular tissue type. Previous studies have demonstrated the presence of three modular regions in the genomes of DWV-like viruses the 5'-UTR, the leader/capsid-encoding region and the region encoding the non-structural proteins, which can be exchanged between related viruses (Moore et al., 2011). No DWV was found in either the synganglion library by RT-PCR, which was made from 50 individuals, or in other mite tissues. This is surprising given that specific primers spanning the recombination point at the capsid / non-structural boundary detected not only the VDV-1 virus, but the recombinant DWV /VDV-1 strain. This is in agreement with the sequencing data from the transcriptome and differs from recombinants found previously (Moore et al., 2011) suggesting that virus recombination and mutation is not homogenous across *Varroa* populations. There is some disagreement as to whether DWV or VDV-1 replicates in *Varroa* (Ongus et al., 2004), (Yue and Genersch, 2005), (Shah et al., 2009) but using strand-specific qPCR we were able to demonstrate that negative VDV-1 strands were present in the synganglia, gut, Malpighian tissue and whole mites. This widespread replication could have implications in transmission of high titres of pathogenic virus to honey bee pupa.

The presence of positive and negative strand VDV-1 in the synganglia of *Varroa* could be significant in its transmission cycle. Thogoto virus in ticks has previously been shown to replicate in the tick synganglia where it is transmitted along neural tissue directly to the salivary glands for transmission to the host via salivary secretions (Kaufman et al., 2001, Kaufman and Nuttall, 1999). A similar mode of transmission of VDV-1 from *Varroa* to the honey bee may occur. Termed the "Host Manipulation Hypothesis", it is well

documented that pathogenic and parasitic organisms change the arthropod vector behaviour to enhance transmission, including for arthropod-borne viruses (Ingwell et al., 2012). For example, tick-borne encephalitis virus alters the tick vector behaviour to exhibit more aggressive and risky behaviour which favours transmission of the virus to the mammalian host (Belova et al., 2012). Replication of VDV-1 in the *Varroa* synganglion, as demonstrated in the present study, could similarly alter *Varroa* behaviour to favour transmission.

A wide range of neuropeptides from arthropods have been characterized but the majority are from the Hexapoda and Crustacea with very few from mites and ticks (Christie et al., 2011). Genetic analysis estimates that acari diverged from insects and crustacean 750 million years ago (Weinstock et al., 2006) and *Varroa* from ticks over 300 million years ago (Cornman et al., 2010) and so it does not necessarily follow that neuropeptides from acari have modes of action similar to either insects and crustacean. Most chelicerate neuropeptides characterized come from just two species of tick, *I. scapularis* (Neupert et al., 2009, Neupert et al., 2005, Simo et al., 2013, Veenstra, 2009) and *Dermacentor variabilis* (Bissinger et al., 2011). We present here a CHH-like peptide with motifs and conserved cysteines in common with previously characterized members of the CHH-superfamily that is expressed in both the synganglion and in isolated gut tissue and Malpighian tubules. It is unknown whether our preparations of gut tissue that showed presence of mite CHH-like peptide by RT-PCR were picking up transcripts from neurons innervating the gut or from gut tissue *per se*. In crustaceans, CHH-like mRNA transcripts have been reported in heart, gills, antennal gland and even the spermatophore sac (Webster et al., 2012). In insects, CHH-like mRNAs have not been found outside the central and peripheral nervous system except in *Tribolium castaneum* where transcripts are abundantly expressed in the midgut at the late larval stage (Begum et al., 2009).

Although a member of the CHH-superfamily, the *Varroa* CHH-like peptide is phylogenetically distinct from the currently characterized superfamily subgroups, falling into a clade shared with other acari and this may be reflected in its functional role. To date, no functional characterization of any acarine CHH-like neuropeptide is available (Christie et al., 2011). CHH-superfamily peptides can be split into subgroups that have distinct roles in arthropods (Jia et al., 2012). CHH1 and CHH2 primarily regulate blood glucose metabolism due to environmental conditions (Chung and Zmora, 2011, Chung et al., 2010). Members of

the MIH subfamily inhibit ecdysteroid synthesis and delay moulting in crustaceans (Nakatsuji et al., 2009). The closely related subgroup GIH, also called vitellogenesis-inhibiting hormone (VIH), suppress the production of methyl farnesoate influencing vitellogenesis and somatic and gonadal growth (Wainwright et al., 1996). Ion Transport Protein (ITP) peptides found in locusts has shown that they regulate Cl^- ion mediated water resorption in ileum (Audsley et al., 1992) and it has been proposed that this is similar to the CHH role in osmoregulation during moulting in crabs (Chung et al., 2010), (Carley, 1978). In *Tribolium* beetles RNAi knockdown of ITP variants suggested a role in moulting and in egg production (Begum et al., 2009)

Clearly, the CHH-superfamily neural peptides are multifunctional and exhibit multiple physiological roles. Although phylogenetic analysis determined the L1C01 contig resides within a distinct acarine clade, no inference of function was gleaned from this analysis. The *Varroa* used for construction of the synganglion library in this study were all mature, mated, adult females emerging from brood cells. Such *Varroa* have moulted to the last stage and have fully formed ovaries and have mated, hence, the *Varroa* CHH-like is unlikely to exhibit moulting-inhibiting or gonad-inhibiting activity of MIH and GIH, respectively, in these individuals. However, these *Varroa* do search for new brood cells containing bee larvae where they will lay their eggs, but this oviposition does not take place until such brood cells have been located and certain cues trigger vitellogenesis (Frey et al., 2013). So, it is plausible that the *Varroa* CHH-like neuropeptide identified in the present study exhibits the vitellogenesis-inhibiting activity of a VIH neuropeptide. Using the RNAi gene-knockdown methods described in this paper in a reverse-genetics approach, researchers could ascertain whether the *Varroa* CHH-like neuropeptide is involved in arrested vitellogenesis and oviposition.

Allatostatins (ASTs) are a neural peptide superfamily found across a range of taxa and are pleiotropic in function including being involved in the inhibition of juvenile hormone synthesis in insects (Stay and Tobe, 2007). Three families of ASTs (A-, B-, and C-type) have been identified, to date. EST sequence analysis identified a B-type AST from *Varroa* that is expressed specifically in the synganglion and elicits a lethal phenotype when knocked-down by RNAi. In the chelicerates, ASTs have been recently identified by *in silico* analysis (Christie et al., 2011, Veenstra et al., 2012), but are yet to be functionally characterised. The B-type family of ASTs are C-terminally amidated peptides with tryptophan in the second and

ninth positions, and are termed the W(X)6Wamide ASTs. This family of peptides is multifunctional and function as myoinhibitory peptides but also demonstrate prothoracicostatic activity (Bendena and Tobe, 2012). The *Varroa* B-type AST pro-hormone contains cleavage sites and W(X)6W motifs that suggest each pro-hormone is processed into three distinct hormone peptides that are significantly different in sequence but each retain the W(X)6W motif. Differences in sequence in small hormonal peptides can have huge effects on the efficacy of hormones of different target receptors and it may be that the three predicted peptides cleaved from the pro-hormone could have different functions or distribution. A putative B-type AST from *M. occidentalis* has similar cleavage and peptide structure to *Varroa*, but a B-type AST isolated from *T. urtica* contains six putative hormone peptides. No functional characterisation of chelicerate allatostatins has yet been determined, however, the highly lethal phenotype presented here on adult female *Varroa* demonstrates that it may play a crucial role outside of reproductive and moult physiology. Gene-knockdown of the *Varroa* AST by the non-invasive immersion approach demonstrated in the present study would permit investigations into the functional role of this neuropeptide hormone. Indeed, the demonstration that the expression of genes within the central nervous can be readily suppressed suggests this approach could be applied more broadly to other neuropeptides and in other arthropods.

dsRNAi gene knockdown was first described in *Varroa* on a mu-type glutathione S-transferase (Campbell et al., 2010) demonstrating that gene expression in the adult *Varroa*, can be specifically inhibited by microinjection or the direct soaking of mites with target specific dsRNA. It has been shown in ticks (Karim et al., 2008) and *Drosophila* (Dzitoyeva et al., 2001) that when dsRNA is injected into the hemocoel, the dsRNA can affect the exposed synganglia. We have confirmed this finding in *Varroa* by targeting neuropeptides and demonstrated the effect can be achieved by the less invasive method of dsRNA administration immersion which is also more amenable to higher throughput. The first targets for the direct management of *Varroa* were validated by the knockdown of a cocktail of 14 targets that reduced *Varroa* populations in mini-honey bee colonies by 53% compared to control (Garbian et al., 2012). In this current study we demonstrate the first evidence of direct lethality to individual mites using a single target with dsRNA-AST killing 54% mites after 72 hours compared to controls. Mortality was also seen in mites treated with dsRNA-CHH with ~25% mortality compared to controls after 72h. In addition a repeatable and significant cohort of dsRNA-CHH treated mites displayed a shaking phenotype after 24-48h.

Field studies of RNAi knockdown in *Varroa* targeted a range of 14 housekeeping genes (Garbian et al., 2012) rather than neural peptides, and it may be that increasing the number of neural targets, as shown here, will increase the efficacy of RNAi as a means of control.

In summary, an EST transcriptome library constructed from the synganglia of *Varroa* mites has identified pathogenic honey bee viruses, potential acaricide targets and critical neurohormones (CHH-like and B-type AST). *Varroa destructor* virus was found replicating in the synganglion and it is postulated that this may alter the behaviour of the *Varroa*. AST gene's expression was restricted to the synganglion and we demonstrated that such genes are amenable to gene-knockdown by administration of the appropriate dsRNA by a simple and trauma-free immersion method. Knockdown of the CHH and AST transcript levels demonstrates that future investigation of neurohormonal function by RNAi in mites and, more broadly, other arthropods is possible and also opens the door to the control of *Varroa* and other pest species by targeting critical genes residing in the central and peripheral nervous system.

Acknowledgements

This work was funded by BBSRC-LINK grant # BB/J01009X/1 and Vita Europe Ltd. We are grateful to the Scottish Beekeepers Association, especially Mr Phil McAnespie in supporting this work at its inception. We acknowledge partial funding from a Genesis Faraday SPARK Award, part of a Scottish Government SEEKIT project for the early part of this work. We are grateful to Prof David Evans for his advice on *Varroa destructor* viruses.

Figure 1. Identity and functional grouping of ESTs from synganglia into clusters. (A) BLAST analysis of transcripts classified EST clusters as mitochondrial sequences, putative identity, unknown function but present in the *Varroa* genome database or novel sequences. (B) Transcripts with a putative identity were further divided into functional categories by analysing GO identity and homology to known genes.

Figure 2. *Varroa destructor* synganglia contain pathogenic strains of deformed wing virus known as VDV-1. (A) Pairwise sequence comparison scores of synganglia EST contigs (L1A04 and L4C04) with both DWV and VDV-1. Scores represent percentage of sequence identity between pairs. (B) Detection of positive-strand virus RNA in the synganglia library was determined by an initial PCR using generic DWV primers followed by nested PCR using primers specific to four DWV recombinants. (C) Negative-strand VDV-1 virus was detected in the synganglia and gut by strand specific RT-qPCR representing replicating virus.

Figure 3. Phylogenetic tree of members of the Crustacean CHH/MIH/GIH/VIH/ITP neurohormone superfamily with homology to *Varroa destructor* L1C01. (A) Bootstrap consensus phylogenetic trees place the Acari CHH-like peptides in a separate clade from currently characterized members of the CHH superfamily. (B) Radiation phylogram of CHH superfamily places Acari CHH superfamily on a separate branch. Initial discrimination of sites by GUIDANCE (PRANK) (Penn et al., 2010) resulted in 67 conserved amino acids considered in MEGA 6 analysis (Tamura et al., 2013). Phylogeny was inferred using the Maximum likelihood method. The bootstrap consensus tree inferred from 1000 replicates (Felsenstein, 1985) is taken to represent the evolutionary history of the taxa analyzed. Branches corresponding to partitions reproduced in less than 50% bootstrap replicates are collapsed. The evolutionary distances were computed using the LG + G + I statistical analysis model (Jones et al., 1992) and are in the units of the number of amino acid substitutions per site. All positions containing gaps and missing data were eliminated. There were a total of 51 positions in the final dataset. Accession numbers are as follows; *Cancer pagurus* (CAC05347.1), *Callinectes sapidus* (AAA69029), *Scylla serrate* (AAL99355.1), *Charybdis feriata* (AAC64785.1), *Carcinus maenas* (CAA53591.1), *Cancer pagurus b* (2208452A), *Nephrops norvegicus* (AAK58133.2), *Macrobrachium rosenbergii* (AAL37948.1), *Penaeus japonicas* (P55847.2), *Fenneropenaeus chinensis* (AAL55258.4), *Penaeus monodon* (BAB70610.1), *Penaeus monodon b* (BAB69830.1), *Metapenaeus ensis* (AAC27452.1), *Trachypenaeus curvirostris* (AAL55259.1), *Eriocheir sinensis* (AAL55256.1), *Litopenaeus stylirostris* (AAL55257.1), *Procambarus clarkia* (P55848.1), *Jasus lalandii* (P83220.1), *Armadillidium vulgare* (P83627.1), *Caenorhabditis elegans* (NP_501985.1), *Homarus americanus* (gCAA38611.1), *Procambarus clarkia b* (BAA89003.1), *Procambarus bouvieri* (P55845.1), *Jasus lalandii b* (P56687.1), *Litopenaeus vannamei* (AAN86056.1), *Macrobrachium lanchesteri* (AAC36310.1), *Armadillidium vulgare* (P30814.1), *Libinia emarginata* (P56688.1), *Cancer pagurus* (P81032.1), *Carcinus maenas b* (P14944.2), *Penaeus japonicas b* (P81700.1), *Penaeus monodon b* (AAC84143.1), *Litopenaeus vannamei b* (CAA68067.1), *Marsupenaeus japonicas b* (BAA13481.1), *Ixodes scapularis* (XP_002400720.1), *Varroa destructor* (L1C01), *Metaseiulus occidentalis* (XP_003741125.1), *Locusta migratoria* (AAD20821.1), *Schistocerca gregaria* (AAB16823.1), *Nasonia vitripennis* (XP_001604056.1), *Tribolium castaneum* (NP_001076808.1), *Anopheles gambiae* (EAA09451.4), *Bombyx mori* (Q9NL55.1), *Manduca sexta* (AAY29658.1), *Folsomia candida* (ACF15252.1), *Drosophila melanogaster* (AAL25507.1), *Tetranychus urticae* (tetur06g00590), *Dermacentor variabilis* (ACC99599.1). CHH = Crustacean hyperglycaemic hormone; MIH = moult-inhibiting

hormone; GIH = gonad-inhibiting hormone; VIH = vitelogenesis-inhibiting hormone; ITP = ion transport peptide. *Varroa destructor* CHH shares functional and structural residues with a homologue in *M. occidentalis* (C). Methionine start codon is marked with M above sequence. Sequence highlights represent cleavage sites for Signal peptide (red with asterix above) and pro-peptide (green with arrow above). Blue highlights with C above show the location of 6 canonical cysteines intrinsic to the folding of CHH neuropeptides and yellow highlights the PFAM CHH superfamily motif.

Figure 4. *Varroa destructor* AST is homologous to B-type allatostatins in insects, crustacea and other acari. (A) Mature peptide sequences aligned in Clustal Omega all contain the allatostatin motif W(X)6W and glycine- amidinated N terminii. Conserved tryptophan residues are highlighted. Accession numbers are as follows; *M. occidentalis* (XP_003741714.1), *T. urtica* (tetur10g00930), *I. scapularis* (XP_002434041.1, *T. castaneum* (NP_001137202.1), *A. aegypti* (XP_001655873.1), *D. ponderosae* (ENN80203.1), *C. quinquefasciatus* (XP_001842234.1), *P. humanus* (XP_002422676.1), *C. capitata* (XP_004530893.1), *M. domestica* (XP_005178794.1), *B. mori* (P82003), *G. bimaculatus* (Q5QRY7), *D. pulex* (E9GSK4). (B) Pro-hormone peptide structures vary between species with lysine / arginine cleavage sites resulting in multiple predicted mature peptides. Amino acid length is displayed after pro-peptide structure. SP = signal peptide. Numbered domains = mature peptides.

Figure 5. *Varroa destructor* CHH is distributed in both synganglion, gut and Malpighian tubule whereas AST is restricted to the synganglion. Expression of Vd-CHH and Vd-AST transcripts were determined by RT-PCR in different tissues of *V. destructor* using CHH and AST specific primers and gel loading normalized to actin. Syn = synganglion; Mite - syng = *Varroa* with synganglion removed and MT = Malpighian tubules.

Figure 6. Extent and persistence of dsRNA-VdAST gene knockdown and subsequent mortality in adult *V. destructor*. (A) Semi-quantitative VdAST RT-PCR band intensities were determined by densitometry and the relative abundance to the within-sample actin band calculated. The extent of VdAST : actin knockdown is presented as a percentage for VdAST-dsRNA injected mites and control mites injected with LacZ-dsRNA at each given time point. Data are means \pm SEM (n = 3 individual mites). Asterisks represent significant difference ($P < 0.001$) between treatments determined by Student's t-test. (B) Survival was monitored post-injection with either VdAST or LacZ-dsRNA. Mites were monitored every 12h with dead mites removed from assay chambers. LacZ (4 groups of 11 mites, n=4), VdAST (3 groups of 10 mites, n =3). Data are means of each chamber \pm SEM. Asterisks represent significant differences between treatments (**= $P < 0.01$; *** = $P < 0.001$) determined by Student's t-test.

Figure 7. Extent and persistence of dsRNA-VdCHH gene knockdown and subsequent mortality in adult *V. destructor*. (A) Semi-quantitative VdCHH RT-PCR band intensities were determined by densitometry and the relative abundance to the within-sample actin band calculated. The extent of VdCHH : actin knockdown is presented as a percentage for VdCHH-dsRNA injected mites and control mites injected with LacZ-dsRNA at each given time point. Data are means \pm SEM (n = 3 individual mites). Asterisks represent significant difference ($P < 0.001$) between treatments determined by Student's t-test. (B) Survival was monitored post-injection with either VdCHH or LacZ-dsRNA. Mites were monitored every 12h with dead mites removed from assay chambers. Injected mites assigned to groups of 4, LacZ (3 groups of 7, n = 3), VdCHH (3 groups of 7 mites, n = 3). Data are means of each chamber \pm SEM. Asterisks represent significant differences between treatments ($P < 0.03$).

ACCEPTED MANUSCRIPT

References

- Audsley, N., McIntosh, C., Phillips, J.E., 1992. Isolation of a neuropeptide from locust corpus cardiacum which influences ileal transport. *J. Exp. Biol.* 173, 261-274.
- Begum, K., Li, B., Beeman, R.W., Park, Y., 2009. Functions of ion transport peptide and ion transport peptide-like in the red flour beetle *Tribolium castaneum*. *Insect Biochem. Mol. Biol.* 39, 717-725.
- Belova, O.A., Burenkova, L.A., Karganova, G.G., 2012. Different tick-borne encephalitis virus (TBEV) prevalences in unfed versus partially engorged ixodid ticks - Evidence of virus replication and changes in tick behavior. *Ticks and Tick-Borne Diseases* 3, 240-246.
- Bendena, W.G., Tobe, S.S., 2012. Families of allatoregulator sequences: a 2011 perspective. *Canadian Journal of Zoology-Revue Canadienne De Zoologie* 90, 521-544.
- Bissinger, B.W., Donohue, K.V., Khalil, S.M.S., Grozinger, C.M., Sonenshine, D.E., Zhu, J., Roe, R.M., 2011. Synganglion transcriptome and developmental global gene expression in adult females of the American dog tick, *Dermacentor variabilis* (Acari: Ixodidae). *Insect Mol. Biol.* 20, 465-491.
- Browning, R., Karim, S., 2013. RNA interference-mediated depletion of N-ethylmaleimide sensitive fusion protein and synaptosomal associated protein of 25 kDa results in the inhibition of blood feeding of the Gulf Coast tick, *Amblyomma maculatum*. *Insect Mol. Biol.* 22, 245-257.
- Bryon, A., Wybouw, N., Dermauw, W., Tirry, L., Van Leeuwen, T., 2013. Genome wide gene-expression analysis of facultative reproductive diapause in the two-spotted spider mite *Tetranychus urticae*. *BMC Genomics* 14, 815-815.
- Cabrera, A.R., Donohue, K.V., Roe, R.M., 2009. Regulation of female reproduction in mites: A unifying model for the Acari. *J. Insect Physiol.* 55, 1079-1090.
- Campbell, E.M., Budge, G.E., Bowman, A.S., 2010. Gene-knockdown in the honey bee mite *Varroa destructor* by a non-invasive approach: studies on a glutathione S-transferase. *Parasites & Vectors* 3, 73.
- Carley, W.W., 1978. Water economy of the Earthworm *Lumbricus terrestris* L.: Coping with the terrestrial environment. *J. Exp. Zool.* 205, 71-78.
- Christie, A.E., Nolan, D.H., Ohno, P., Hartline, N., Lenz, P.H., 2011. Identification of chelicerate neuropeptides using bioinformatics of publicly accessible expressed sequence tags. *Gen. Comp. Endocrinol.* 170, 144-155.
- Chung, J.S., Zmora, N., 2011. Multifunctionality of crustacean hyperglycemic hormone neuropeptide family in molting and reproduction of the Blue Crab, *Callinectes Sapidus*. *J. Shellfish Res.* 30, 496-496.

- 753 Chung, J.S., Zmora, N., Katayama, H., Tsutsui, N., 2010. Crustacean hyperglycemic
754 hormone (CHH) neuropeptides family: Functions, titer, and binding to target tissues. *Gen.*
755 *Comp. Endocrinol.* 166, 447-454.
- 756 Cornman, S.R., Schatz, M.C., Johnston, S.J., Chen, Y., Pettis, J., Hunt, G., Bourgeois, L.,
757 Elsik, C., Anderson, D., Grozinger, C.M., Evans, J.D., 2010. Genomic survey of the
758 ectoparasitic mite *Varroa destructor*, a major pest of the honey bee *Apis mellifera*. *BMC*
759 *Genomics* 11, 602.
- 760 Correa-Marques, M.H., de Jong, D., Rosenkranz, P., Goncalves, L.S., 2002. *Varroa*-tolerant
761 Italian honey bees introduced from Brazil were not more efficient in defending themselves
762 against the mite *Varroa destructor* than Carniolan bees in Germany. *Genetics and Molecular*
763 *Research* 1, 153-158.
- 764 de Miranda, J.R., Bailey, L., Ball, B.V., Blanchard, P., Budge, G.E., Chejanovsky, N., Chen,
765 Y., Gauthier, L., Genersch, E., de Graaf, D.C., Ribiere, M., Ryabov, E., De Smet, L., van der
766 Steen, J.J.M., 2013. Standard methods for virus research in *Apis mellifera*. *J. Apic. Res.* 52,
767 UNSP 52.4.22-UNSP 52.4.22.
- 768 Di Prisco, G., Pennacchio, F., Caprio, E., Boncristiani, H.F., Jr., Evans, J.D., Chen, Y., 2011.
769 *Varroa destructor* is an effective vector of Israeli acute paralysis virus in the honeybee, *Apis*
770 *mellifera*. *J. Gen. Virol.* 92, 151-155.
- 771 Dietemann, V., Pflugfelder, J., Anderson, D., Charriere, J., Chejanovsky, N., Dainat, B., de
772 Miranda, J., Delaplane, K., Dillier, F., Fuch, S., Gallmann, P., Gauthier, L., Imdorf, A.,
773 Koeniger, N., Kralj, J., Meikle, W., Pettis, J., Rosenkranz, P., Sammartaro, D., Smith, D.,
774 Yanez, O., Neumann, P., 2012. *Varroa destructor*: research avenues towards sustainable
775 control. *J. Apic. Res.* 51, 125-132.
- 776 Dzitoyeva, S., Dimitrijevic, N., Manev, H., 2001. Intra-abdominal injection of double-
777 stranded RNA into anesthetized adult *Drosophila* triggers RNA interference in the central
778 nervous system. *Mol. Psychiatry* 6, 665-670.
- 779 Felsenstein, J., 1985. Confidence-limits on phylogenies - an approach using the bootstrap.
780 *Evolution* 39, 783-791.
- 781 Frey, E., Odemer, R., Blum, T., Rosenkranz, P., 2013. Activation and interruption of the
782 reproduction of *Varroa destructor* is triggered by host signals (*Apis mellifera*). *J. Invertebr.*
783 *Pathol.* 113, 56-62.
- 784 Friis, K., Damgaard, C., Holmstrup, M., 2004. Sublethal soil copper concentrations increase
785 mortality in the earthworm *Aporrectodea caliginosa* during drought. *Ecotoxicol. Environ.*
786 *Saf.* 57, 65-73.
- 787 Gallai, N., Salles, J., Settele, J., Vaissiere, B.E., 2009. Economic valuation of the
788 vulnerability of world agriculture confronted with pollinator decline. *Ecol. Econ.* 68, 810-
789 821.

- 790 Garbian, Y., Maori, E., Kalev, H., Shafir, S., Sela, I., 2012. Bidirectional transfer of RNAi
791 between honey bee and *Varroa destructor*: Varroa gene silencing reduces Varroa population.
792 Plos Pathogens 8, e1003035-e1003035.
- 793 Genersch, E., Aubert, M., 2010. Emerging and re-emerging viruses of the honey bee (*Apis*
794 *mellifera* L.). Vet. Res. 41, 54.
- 795 Highfield, A.C., El Nagar, A., Mackinder, L.C.M., Noel, L.M.-J., Hall, M.J., Martin, S.J.,
796 Schroeder, D.C., 2009. Deformed wing virus implicated in overwintering Honeybee colony
797 losses. Appl. Environ. Microbiol. 75, 7212-7220.
- 798 Hoy, M.A., Yu, F., Meyer, J.M., Tarazona, O.A., Jeyaparakash, A., Wu, K., 2013.
799 Transcriptome sequencing and annotation of the predatory mite *Metaseiulus occidentalis*
800 (Acari: Phytoseiidae): a cautionary tale about possible contamination by prey sequences. Exp.
801 Appl. Acarol. 59, 283-296.
- 802 Ingwell, L.L., Eigenbrode, S.D., Bosque-Perez, N.A., 2012. Plant viruses alter insect
803 behavior to enhance their spread. Scientific Reports 2, 578.
- 804 Jia, C., Hui, L., Cao, W., Lietz, C.B., Jiang, X., Chen, R., Catherman, A.D., Thomas, P.M.,
805 Ge, Y., Kelleher, N.L., Li, L., 2012. High-definition *de novo* sequencing of crustacean
806 hyperglycemic hormone (CHH)-family neuropeptides. Molecular & Cellular Proteomics 11,
807 1951-1964.
- 808 Jones, D.T., Taylor, W.R., Thornton, J.M., 1992. The rapid generation of mutation data
809 matrices from protein sequences. Computer Applications in the Biosciences 8, 275-282.
- 810 Karim, S., Kenny, B., Troiano, E., Mather, T.N., 2008. RNAi-mediated gene silencing in tick
811 synganglia: A proof of concept study. BMC Biotechnology 8, 30-30.
- 812 Kaufman, W.R., Bowman, A.S., Nuttall, P.A., 2001. Salivary fluid secretion in the ixodid
813 tick *Rhipicephalus appendiculatus* is inhibited by Thogoto virus infection. Experimental and
814 Applied Acarology 25, 661-674.
- 815 Kaufman, W.R., Nuttall, P.A., 1999. Secretion of Thogoto virus in saliva of the Ixodid tick,
816 *Amblyomma Variegatum*. Acarology IX, 2.
- 817 Khila, A., Grbic, M., 2007. Gene silencing in the spider mite *Tetranychus urticae*: dsRNA
818 and siRNA parental silencing of the Distal-less gene. Dev. Genes Evol. 217, 241-251.
- 819 Kleijn, D., Winfree, R., Bartomeus, I., Carvalheiro, L.G., Henry, M., Isaacs, R., Klein, A.,
820 Kremen, C., M'Gonigle, L.K., Rader, R., Ricketts, T.H., Williams, N.M., Lee Adamson, N.,
821 Ascher, J.S., Baldi, A., Batary, P., Benjamin, F., Biesmeijer, J.C., Blitzer, E.J., Bommarco,
822 R., Brand, M.R., Bretagnolle, V., Button, L., Cariveau, D.P., Chifflet, R., Colville, J.F.,
823 Danforth, B.N., Elle, E., Garratt, M.P.D., Herzog, F., Holzschuh, A., Howlett, B.G., Jauker,
824 F., Jha, S., Knop, E., Krewenka, K.M., Le Feon, V., Mandelik, Y., May, E.A., Park, M.G.,
825 Pisanty, G., Reemer, M., Riedinger, V., Rollin, O., Rundlof, M., Sardinas, H.S., Scheper, J.,
826 Sciligo, A.R., Smith, H.G., Steffan-Dewenter, I., Thorp, R., Tschamntke, T., Verhulst, J.,
827 Viana, B.F., Vaissiere, B.E., Veldtman, R., Westphal, C., Potts, S.G., 2015. Delivery of crop


- 828 pollination services is an insufficient argument for wild pollinator conservation. *Nature*
829 *communications* 6, 7414-7414.
- 830 Krogh, A., Larsson, B., von Heijne, G., Sonnhammer, E.L.L., 2001. Predicting
831 transmembrane protein topology with a hidden Markov model: Application to complete
832 genomes. *J. Mol. Biol.* 305, 567-580.
- 833 Kwon, T.H., Hager, H., Nejsum, L.N., Andersen, M.L.E., Frokiaer, J., Nielsen, S., 2001.
834 Physiology and pathophysiology of renal aquaporins. *Semin. Nephrol.* 21, 231-238.
- 835 Lees, K., Woods, D.J., Bowman, A.S., 2010. Transcriptome analysis of the synganglion from
836 the brown dog tick, *Rhipicephalus sanguineus*. *Insect Mol. Biol.* 19, 273-282.
- 837 Lees, K., Bowman, A.S., 2007. Tick neurobiology: recent advances and the post-genomic
838 era. *Invertebr. Neurosci.* 7, 183-198.
- 839 Martin, S., 1998. A population model for the ectoparasitic mite *Varroa jacobsoni* in honey
840 bee (*Apis mellifera*) colonies. *Ecol. Model.* 109, 267-281.
- 841 Martin, S.J., Ball, B.V., Carreck, N.L., 2013. The role of deformed wing virus in the initial
842 collapse of varroa infested honey bee colonies in the UK. *J. Apic. Res.* 52, UNSP 52.5.12.
- 843 Martin, S.J., Highfield, A.C., Brettell, L., Villalobos, E.M., Budge, G.E., Powell, M.,
844 Nikaido, S., Schroeder, D.C., 2012. Global Honey bee viral landscape altered by a parasitic
845 mite. *Science* 336, 1304-1306.
- 846 Moore, J., Jironkin, A., Chandler, D., Burroughs, N., Evans, D.J., Ryabov, E.V., 2011.
847 Recombinants between Deformed wing virus and *Varroa destructor* virus-1 may prevail in
848 *Varroa destructor*-infested honeybee colonies. *J. Gen. Virol.* 92, 156-161.
- 849 Nakatsuji, T., Lee, C., Watson, R.D., 2009. Crustacean molt-inhibiting hormone: Structure,
850 function, and cellular mode of action. *Comparative Biochemistry and Physiology A-*
851 *Molecular & Integrative Physiology* 152, 139-148.
- 852 Neupert, S., Predel, R., Russell, W.K., Davies, R., Pietrantonio, P.V., Nachman, R.J., 2005.
853 Identification of tick periviscerokinin, the first neurohormone of Ixodidae: Single cell
854 analysis by means of MALDI-TOF/TOF mass spectrometry. *Biochem. Biophys. Res.*
855 *Commun.* 338, 1860-1864.
- 856 Neupert, S., Russell, W.K., Predel, R., Russell, D.H., Strey, O.F., Teel, P.D., Nachman, R.J.,
857 2009. The neuropeptidomics of *Ixodes scapularis* synganglion. *Journal of Proteomics* 72,
858 1040-1045.
- 859 Ongus, J.R., Peters, D., Bonmatin, J., Bengsch, E., Vlak, J.M., van Oers, M.M., 2004.
860 Complete sequence of a picorna-like virus of the genus Iflavirus replicating in the mite
861 *Varroa destructor*. *J. Gen. Virol.* 2004, 3747-3755.
- 862 Penn, O., Privman, E., Ashkenazy, H., Landan, G., Graur, D., Pupko, T., 2010. GUIDANCE:
863 a web server for assessing alignment confidence scores. *Nucleic Acids Res.* 38, W23-W28.

- 864 Price, D.R.G., Gatehouse, J.A., 2008. RNAi-mediated crop protection against insects. Trends
865 Biotechnol. 26, 393-400.
- 866 Rodriguez-Dehaibes, S.R., Otero-Colina, G., Villanueva-Jimenez, J.A., Corcuera, P., 2011.
867 Susceptibility of *Varroa destructor* (Gamasida: Varroidae) to four pesticides used in three
868 Mexican apicultural regions under two different management systems. Int. J. Acarol. 37, 441-
869 447.
- 870 Rosenkranz, P., Aumeier, P., Ziegelmann, B., 2010. Biology and control of *Varroa*
871 *destructor*. J. Invertebr. Pathol. 103, S96-S119.
- 872 Ryabov, E.V., Wood, G.R., Fannon, J.M., Moore, J.D., Bull, J.C., Chandler, D., Mead, A.,
873 Burroughs, N., Evans, D.J., 2014. A virulent strain of Deformed wing virus (DWV) of
874 Honeybees (*Apis mellifera*) prevails after *Varroa destructor*-mediated, or *in vitro*,
875 Transmission. Plos Pathogens 10, e1004230.
- 876 Schwarz, A., von Reumont, B.M., Erhart, J., Chagas, A.C., Ribeiro, J.M.C., Kotsyfakis, M.,
877 2013. *De novo* *Ixodes ricinus* salivary gland transcriptome analysis using two next-generation
878 sequencing methodologies. FASEB journal : official publication of the Federation of
879 American Societies for Experimental Biology 27, 4745-56.
- 880 Scott, J.G., Michel, K., Bartholomay, L.C., Siegfried, B.D., Hunter, W.B., Smagghe, G., Zhu,
881 K.Y., Douglas, A.E., 2013. Towards the elements of successful insect RNAi. J. Insect
882 Physiol. 59, 1212-1221.
- 883 Shah, K.S., Evans, E.C., Pizzorno, M.C., 2009. Localization of Deformed wing virus (DWV)
884 in the brains of the honeybee, *Apis mellifera* Linnaeus. Virology Journal 6, 182-182.
- 885 Simo, L., Koci, J., Park, Y., 2013. Receptors for the neuropeptides, myoinhibitory peptide
886 and SIFamide, in control of the salivary glands of the blacklegged tick *Ixodes scapularis*.
887 Insect Biochem. Mol. Biol. 43, 376-387.
- 888 Stay, B., Tobe, S.S., 2007. The role of allatostatins in juvenile hormone synthesis in insects
889 and crustaceans. Annu. Rev. Entomol. 52, 277-299.
- 890 Tamura, K., Stecher, G., Peterson, D., Filipski, A., Kumar, S., 2013. MEGA6: Molecular
891 Evolutionary Genetics Analysis version 6.0. Mol. Biol. Evol. 30, 2725-2729.
- 892 vanEngelsdorp, D., Evans, J.D., Saegerman, C., Mullin, C., Haubruge, E., Nguyen, B.K.,
893 Frazier, M., Frazier, J., Cox-Foster, D., Chen, Y., Underwood, R., Tarpy, D.R., Pettis, J.S.,
894 2009. Colony Collapse Disorder: A Descriptive Study. Plos One 4, e6481.
- 895 vanEngelsdorp, D., Meixner, M.D., 2010. A historical review of managed honey bee
896 populations in Europe and the United States and the factors that may affect them. J. Invertebr.
897 Pathol. 103, S80-S95.
- 898 Veenstra, J.A., 2000. Mono- and dibasic proteolytic cleavage sites in insect neuroendocrine
899 peptide precursors. Arch. Insect Biochem. Physiol. 43, 49-63.

- 900 Veenstra, J.A., 2009. Allatostatin C and its paralog allatostatin double C: The arthropod
901 somatostatins. *Insect Biochem. Mol. Biol.* 39, 161-170.
- 902 Veenstra, J.A., Rombauts, S., Grbic, M., 2012. *In silico* cloning of genes encoding
903 neuropeptides, neurohormones and their putative G-protein coupled receptors in a spider
904 mite. *Insect Biochem. Mol. Biol.* 42, 277-295.
- 905 Wainwright, G., Webster, S.G., Wilkinson, M.C., Chung, J.S., Rees, H.H., 1996. Structure
906 and significance of mandibular organ-inhibiting hormone in the crab, *Cancer pagurus* -
907 Involvement in multihormonal regulation of growth and reproduction. *J. Biol. Chem.* 271,
908 12749-12754.
- 909 Wang, H., Xie, J., Shreeve, T.G., Ma, J., Pallett, D.W., King, L.A., Possee, R.D., 2013.
910 Sequence recombination and conservation of *Varroa destructor* Virus-1 and Deformed wing
911 virus in field collected Honey bees (*Apis mellifera*). *Plos One* 8, UNSP e74508.
- 912 Wang, R.W., Huang, Z.Y., Dong, K., 2003. Molecular characterization of an arachnid sodium
913 channel gene from the varroa mite (*Varroa destructor*). *Insect Biochem. Mol. Biol.* 33, 733-
914 739.
- 915 Webster, S.G., Keller, R., Dirksen, H., 2012. The CHH-superfamily of multifunctional
916 peptide hormones controlling crustacean metabolism, osmoregulation, moulting, and
917 reproduction. *Gen. Comp. Endocrinol.* 175, 217-233.
- 918 Weinstock, G.M., Robinson, G.E., Gibbs, R.A., Worley, K.C., Evans, J.D., Maleszka, R.,
919 Robertson, H.M., Weaver, D.B., Beye, M., Bork, P., Elsik, C.G., Hartfelder, K., Hunt, G.J.,
920 Zdobnov, E.M., Amdam, G.V., Bitondi, M.M.G., Collins, A.M., Cristino, A.S., Lattorff,
921 H.M.G., Lobo, C.H., Moritz, R.F.A., Nunes, F.M.F., Page, R.E., Jr., Simoes, Z.L.P., Wheeler,
922 D., Carninci, P., Fukuda, S., Hayashizaki, Y., Kai, C., Kawai, J., Sakazume, N., Sasaki, D.,
923 Tagami, M., Albert, S., Baggerman, G., Beggs, K.T., Bloch, G., Cazzamali, G., Cohen, M.,
924 Drapeau, M.D., Eisenhardt, D., Emore, C., Ewing, M.A., Fahrbach, S.E., Foret, S.,
925 Grimmelikhuijzen, C.J.P., Hauser, F., Hummon, A.B., Huybrechts, J., Jones, A.K.,
926 Kadowaki, T., Kaplan, N., Kucharski, R., Leboulle, G., Linial, M., Littleton, J.T., Mercer,
927 A.R., Richmond, T.A., Rodriguez-Zas, S.L., Rubin, E.B., Sattelle, D.B., Schlupalius, D.,
928 Schoofs, L., Shemesh, Y., Sweedler, J.V., Velarde, R., Verleyen, P., Vierstraete, E.,
929 Williamson, M.R., Ament, S.A., Brown, S.J., Corona, M., Dearden, P.K., Dunn, W.A.,
930 Elekonich, M.M., Fujiyuki, T., Gattermeier, I., Gempe, T., Hasselmann, M., Kadowaki, T.,
931 Kage, E., Kamikouchi, A., Kubo, T., Kucharski, R., Kunieda, T., Lorenzen, M.D., Milshina,
932 N.V., Morioka, M., Ohashi, K., Overbeek, R., Ross, C.A., Schioett, M., Shippy, T., Takeuchi,
933 H., Toth, A.L., Willis, J.H., Wilson, M.J., Gordon, K.H.J., Letunic, I., Hackett, K., Peterson,
934 J., Felsenfeld, A., Guyer, M., Solignac, M., Agarwala, R., Cornuet, J.M., Monnerot, M.,
935 Mougél, F., Reese, J.T., Vautrin, D., Gillespie, J.J., Cannone, J.J., Gutell, R.R., Johnston,
936 J.S., Eisen, M.B., Iyer, V.N., Iyer, V., Kosarev, P., Mackey, A.J., Solovyev, V., Souvorov,
937 A., Aronstein, K.A., Bilikova, K., Chen, Y.P., Clark, A.G., Decanini, L.I., Gelbart, W.M.,
938 Hetru, C., Hultmark, D., Imler, J., Jiang, H., Kanost, M., Kimura, K., Lazzaro, B.P., Lopez,
939 D.L., Simuth, J., Thompson, G.J., Zou, Z., De Jong, P., Sodergren, E., Csuroes, M.,
940 Milosavljevic, A., Osoegawa, K., Richards, S., Shu, C., Duret, L., Elhaik, E., Graur, D.,
941 Anzola, J.M., Campbell, K.S., Childs, K.L., Collinge, D., Crosby, M.A., Dickens, C.M.,
942 Grametes, L.S., Grozinger, C.M., Jones, P.L., Jorda, M., Ling, X., Matthews, B.B., Miller, J.,
943 Mizzen, C., Peinado, M.A., Reid, J.G., Russo, S.M., Schroeder, A.J., St Pierre, S.E., Wang,

- 944 Y., Zhou, P., Jiang, H., Kitts, P., Ruef, B., Venkatraman, A., Zhang, L., Aquino-Perez, G.,
 945 Whitfield, C.W., Behura, S.K., Berlocher, S.H., Sheppard, W.S., Smith, D.R., Suarez, A.V.,
 946 Tsutsui, N.D., Wei, X., Wheeler, D., Havlak, P., Li, B., Liu, Y., Sodergren, E., Jolivet, A.,
 947 Lee, S., Nazareth, L.V., Pu, L., Thorn, R., Stolc, V., Newman, T., Samanta, M., Tongprasit,
 948 W.A., Claudianos, C., Berenbaum, M.R., Biswas, S., de Graaf, D.C., Feyereisen, R., Johnson,
 949 R.M., Oakeshott, J.G., Ranson, H., Schuler, M.A., Muzny, D., Chacko, J., Davis, C., Dinh,
 950 H., Gill, R., Hernandez, J., Hines, S., Hume, J., Jackson, L., Kovar, C., Lewis, L., Miner, G.,
 951 Morgan, M., Nguyen, N., Okwuonu, G., Paul, H., Santibanez, J., Savery, G., Svatek, A.,
 952 Villasana, D., Wright, R., Honeybee genome sequencing consort, 2006. Insights into social
 953 insects from the genome of the honeybee *Apis mellifera*. Nature (London) 443, 931-949.
- 954 Wolstenholme, A.J., Bowman, A.S., Sattelle, D.B., 2007. Frontiers in parasite neurobiology:
 955 parasite genomics, neural signalling and new targets for control. Invertebr. Neurosci. 7, 179-
 956 181.
- 957 Yue, C., Genersch, E., 2005. RT-PCR analysis of Deformed wing virus in honeybees (*Apis*
 958 *mellifera*) and mites (*Varroa destructor*). J. Gen. Virol. 86, 3419-3424.

959






Figure 1. Identity and functional classification of clusters.





Figure 2. Varroa destructor virus was present and replicating in the synganglion of *V. destructor*.


A.

<i>V. destructor</i>	-PQPQWNELSGYWG
<i>M. occidentalis</i>	-PQPQWNELSGYWG
<i>T. urtica</i>	-DSPHWNNLRGMWG
<i>I. scapularis</i>	--ENHWNDLSGYWG
<i>A. aegypti</i>	--SEKWNLSSSWG
<i>D. ponderosae</i>	--QPAWNNLKGLWG
<i>C. quinquefasciatus</i>	--SEKWNLSSSWG
<i>P. humanus</i>	SVDSKWAKLQGGWG
<i>C. capitata</i>	S-SKDWAKLHGGWG
<i>M. domesticus</i>	---ATWQKLHGGWG
<i>T. castaneum</i>	---SKWDNFRGSWG
<i>B. mori</i>	----AWSALHGTWG
<i>G. bimaculatus</i>	----AWNNLGSAWG
<i>D. pulex</i>	----SWTQLHGVWG

B.

</





Highlights:

- The first transcriptome of *Varroa destructor*
- Deformed wing virus variant (VDV-1) present and replicating in *Varroa* synganglion (“brain”)
- B-type allatostatin (*VdAST*) and a crustacean hyperglycaemic hormone (*VdCHH*) identified in *Varroa* synganglia
- *VdAST* and *VdCHH* expression knocked down by dsRNAi resulting in significant mortality